

NOAA Technical Report NMFS SSRF-694



Semiclosed Seawater System With Automatic Salinity, Temperature, and Turbidity Control

SID KORN

SEATTLE, WA
September 1975

noaa

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION

National Marine
Fisheries Service

NOAA TECHNICAL REPORTS

National Marine Fisheries Service, Special Scientific Report—Fisheries Series

The major responsibilities of the National Marine Fisheries Service (NMFS) are to monitor and assess the abundance and geographic distribution of fishery resources, to understand and predict fluctuations in the quantity and distribution of these resources, and to establish levels for optimum use of the resources. NMFS is also charged with the development and implementation of policies for managing national fishing grounds, development and enforcement of domestic fisheries regulations, surveillance of foreign fishing off United States coastal waters, and the development and enforcement of international fishery agreements and policies. NMFS also assists the fishing industry through marketing service and economic analysis programs, and mortgage insurance and vessel construction subsidies. It collects, analyzes, and publishes statistics on various phases of the industry.

The Special Scientific Report—Fisheries series was established in 1949. The series carries reports on scientific investigations that document long term continuing programs of NMFS, or intensive scientific reports on studies of restricted scope. The reports may deal with applied fishery problems. The series is also used as a medium for the publication of bibliographies of a specialized scientific nature.

NOAA Technical Reports—NMFS SSRF are available free in limited numbers to governmental agencies, both Federal and State. They are also available in exchange for other scientific and technical publications in the marine sciences. Individual copies may be obtained (unless otherwise noted) from D83, Technical Information Division, Environmental Science Information Center, NOAA, Washington, D.C. 20235. Recent SSRF's are:

619. Macrozooplankton and small nekton in the coastal waters off Vancouver Island (Canada) and Washington, Spring and fall of 1963. By Donald S. Day, January 1971, iii + 94 p., 19 figs., 13 tables.

620. The Trade Wind Zone Oceanography Pilot Study Part IX. The sea level wind field and wind stress values, July 1963 to June 1965. By Gunter R. Seckel, June 1970, iii + 66 p., 5 figs.

621. Predation by sculpins on fall chinook salmon, *Oncorhynchus tshawytscha*, fry of hatchery origin. By Benjamin G. Patten, February 1971, iii + 14 p., 6 figs., 9 tables.

622. Number and lengths, by season, of fishes caught with an otter trawl near Woods Hole, Massachusetts, September 1961 to December 1962. By F. E. Lux and F. E. Nieby, February 1971, iii + 15 p., 3 figs., 19 tables.

623. Apparent abundance, distribution, and migrations of albacore, *Thunnus alalunga*, on the North Pacific longline grounds. By Brian J. Rothschild and Marian Y. Yung, September 1970, v + 37 p., 19 figs., 5 tables.

624. Influence of mechanical processing on the quality and yield of bay scallop meats. By N. B. Webb and F. B. Thomas, April 1971, iii + 11 p., 9 figs., 3 tables.

625. Distribution of salmon and related oceanographic features in the North Pacific Ocean, spring 1968. By Robert B. French, Richard G. Bakkala, Masanao Osako, and Jun Ito, March 1971, iii + 22 p., 19 figs., 3 tables.

626. Commercial fishery and biology of the freshwater shrimp, *Macrobrachium*, in the Lower St. Paul River, Liberia, 1952-53. By George C. Miller, February 1971, iii + 13 p., 8 figs., 7 tables.

627. Calico scallops of the Southeastern United States, 1959-69. By Robert Cummins, Jr., June 1971, iii + 22 p., 23 figs., 3 tables.

628. Fur Seal Investigations—1969. By NMFS, Marine Mammal Biological Laboratory, August 1971, 82 p., 20 figs., 14 tables, 23 appendix A tables, 10 appendix B tables.

629. Analysis of the operations of seven Hawaiian skipjack tuna fishing vessels, June-August 1967. By Richard N. Uchida and Ray F. Sumida, March 1971, v + 25 p., 14 figs., 21 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

630. Blue crab meat I. Preservation by freezing. July 1971, iii + 13 p., 5 figs., 2 tables. II. Effect of chemical treatments on acceptability. By Jurgen H. Strasser, Jean S. Lennon, and Frederick J. King, July 1971, iii + 12 p., 1 fig., 9 tables.

631. Occurrence of thiaminase in some common aquatic animals of the United States and Canada. By R. A. Greig and R. H. Gnaedinger, July 1971, iii + 7 p., 2 tables.

632. An annotated bibliography of attempts to rear the larvae of marine fishes in the laboratory. By Robert C. May, August 1971, iii + 24 p., 1 appendix I table, 1 appendix II table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

633. Blueing of processed crab meat. II. Identification of some factors involved in the blue discoloration of canned crab meat *Callinectes sapidus*. By Melvin E. Waters, May 1971, iii + 7 p., 1 fig., 3 tables.

634. Age composition, weight, length, and sex of herring, *Clupea pallasi*, used for reduction in Alaska, 1929-66. By Gerald M. Reid, July 1971, iii + 25 p., 4 figs., 18 tables.

635. A bibliography of the blackfin tuna, *Thunnus atlanticus* (Lesson). By Grant L. Beardsley and David C. Simmons, August 1971, 10 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

636. Oil pollution on Wake Island from the tanker *R. C. Stoner*. By Reginald M. Gooding, May 1971, iii + 12 p., 8 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

637. Occurrence of larval, juvenile, and mature crabs in the vicinity of Beaufort Inlet, North Carolina. By Donnie L. Dudley and Mayo H. Judy, August 1971, iii + 10 p., 1 fig., 5 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

638. Length-weight relations of haddock from commercial landings in New England, 1931-55. By Bradford E. Brown and Richard C. Hennemuth, August 1971, v + 13 p., 16 figs., 6 tables, 10 appendix A tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

639. A hydrographic survey of the Galveston Bay system, Texas 1963-66. By E. J. Pullen, W. L. Trent, and G. B. Adams, October 1971, v + 13 p., 15 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

640. Annotated bibliography on the fishing industry and biology of the blue crab, *Callinectes sapidus*. By Marlin E. Tagatz and Ann Bowman Hall, August 1971, 94 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

641. Use of tureadfin shad, *Dorosoma petenense*, as live bait during experimental pole and line fishing for skipjack tuna, *Katsuwonus pelamis*, in Hawaii. By Robert T. B. Iversen, August 1971, iii + 10 p., 3 figs., 7 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

642. Atlantic menhaden *Brevoortia tyrannus* resource and fishery—analysis of decline. By Kenneth A. Henry, August 1971, v + 32 p., 40 figs., 5 appendix figs., 3 tables, 2 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

643. Surface winds of the southeastern tropical Atlantic Ocean. By John M. Steigner and Merton C. Ingham, October 1971, iii + 20 p., 17 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

644. Inhibition of flesh browning and skin color fading in frozen fillets of yelloweye snapper (*Lutjanus vivanus*). By Harold C. Thompson, Jr., and Mary H. Thompson, February 1972, iii + 6 p., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

645. Traveling screen for removal of debris from rivers. By Daniel W. Bates, Ernest W. Murphy, and Martin G. Beam, October 1971, iii + 6 p., 6 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

646. Dissolved nitrogen concentrations in the Columbia and Snake Rivers in 1970 and their effect on chinook salmon and steelhead trout. By Wesley J. Ebel, August 1971, iii + 7 p., 2 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

647. Revised annotated list of parasites from sea mammals caught off the west coast of North America. By L. Margolis and M. D. Dailey, March 1972, iii + 23 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.

NOAA Technical Report NMFS SSRF-694

Semiclosed Seawater System With Automatic Salinity, Temperature, and Turbidity Control

SID KORN

SEATTLE, WA
September 1975

5 PP

UNITED STATES
DEPARTMENT OF COMMERCE
Rogers C. B. Morton, Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Robert M. White, Administrator

National Marine
Fisheries Service
Robert W. Schoning, Director



The National Marine Fisheries Service (NMFS) does not approve, recommend or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

CONTENTS

	Page
Introduction	1
Flow description	1
Temperature control	3
Salinity control	3
Alarm system	3
Performance and maintenance	3
Acknowledgments	5
Literature cited	5

Figures

1. Schematic of Tiburon seawater system	2
2. Intake pumps, showing cleanouts and dual piping	2
3. Reservoir, aerator, brine tank, and pneumatic salt delivery line	2
4. Part of main wet laboratory	3
5. Recirculating line in trench	3
6. Schematic of salinity control system	4
7. Control and alarm panel	4

Table

1. Comparison of raw and recirculated water in seawater system	5
--	---

Semiclosed Seawater System With Automatic Salinity, Temperature, and Turbidity Control

SID KORN¹

ABSTRACT

The new seawater system at the Southwest Fishery Center, Tiburon Laboratory, is described. The system delivers up to 450 l/min of filtered, ultraviolet sterilized, temperature- and salinity-controlled seawater suitable for extended holding of marine fish and invertebrates. Unique aspects of the system including provisions for open and/or closed circulation, and the pneumatic salinity control components are described in detail. The design of this facility may offer ideas to others desiring near-oceanic quality seawater from marginal sources.

INTRODUCTION

The purpose of this paper is to describe the new seawater facility of the National Marine Fisheries Service (NMFS), Southwest Fisheries Center, Tiburon Laboratory. Special emphasis is placed on the salinity control components and design features that allow for either open or closed circulation. I hope this information will assist other facilities with water supplies varying in temperature, salinity, and turbidity, in their effort to obtain near-oceanic quality water.

This facility is located on San Francisco Bay where water conditions vary considerably in temperature, salinity, and turbidity, partially caused by runoff from the Sacramento-San Joaquin rivers. The former seawater system, operating from 1967 to 1971, did not control temperature or salinity and had a limited filtering capacity. Fouling of waterlines and tanks, wide fluctuations in temperature (range 9°-20°C) and salinity (range 2-30‰), and bacterial disease problems made research impossible during certain times of the year. This facility is used for physiological research and needs a system that supplies water of near-oceanic quality with constant physical conditions.

Final specifications and design work were completed by consulting engineers. The system was constructed from November 1972 to April 1973.

The new seawater system (Fig. 1) is a semiclosed system which can circulate 450 l/min of filtered, sterilized, temperature- and salinity-controlled water. Most of the system is duplicated to facilitate repairs and cleaning without disruption of the waterflow.

Due to the high cost of salinity and temperature control, most of the water is recirculated. However, some open circulation is required when toxic effluents from experiments are involved and to insure that problems associated with recirculating water do not occur. After water has been recirculated for extended periods through tanks containing a biomass, nitrates accumulate, trace

elements may be lost, and pH may change. Therefore, we always maintain at least 10% of the total flow to open circulation. Toxic effluents are filtered through activated charcoal before being discharged.

FLOW DESCRIPTION

Seawater is pumped by one of two centrifugal pumps (Ace 3 horsepower, hard rubber [Fig. 2]), through a two-stage, automatically backflushing, high-rate sand filter (Baker Model HRB-30). From the filter, the flow is directed to either the sump tank or back into the bay by an electrically actuated valve, which is controlled by a level indicator in the reservoir (Fig. 1). This is done to prevent excess water from entering the system. Dual 3-inch PVC (polyvinyl chloride) lines allow for cleaning of lines and changing of the pump.

The 600-liter epoxy-lined concrete sump tank collects new filtered water and water that has been recirculated through the laboratories. As the water level rises, one of two 10-horsepower centrifugal pumps (Allis Chalmers Model F-4, 316 stainless steel) is activated by a float valve. Water is pumped through a second single-stage sand filter (Baker Model HRB-36) to the 53,000-liter reservoir (Fig. 3). The water enters the reservoir through a redwood splash tray aerator patterned after a similar design at the Southwest Fisheries Center, La Jolla, Calif. (Lasker and Vlymen 1969).

The concrete reservoir is epoxy-lined and is divided by a concrete wall into two sections. Water can be valved to enter and leave one or both sides. This allows for the cleaning of one half of the reservoir without disrupting the flow through the system. The capacity of the tank allows for a 2-h full flow reserve in the event of a failure. By reducing flow rates to 112 l/min this reserve can be extended to 8 h.

Water flows (up to 450 l/min) by gravity from the reservoir to the wet laboratories passing through: an ultraviolet (UV) sterilizer; a heat exchanger; and temperature- and salinity-measuring probes. The flow is then directed into tanks and aquaria as needed (Fig. 4). Depending on the need for open or closed circulation, the effluent line is valved to direct flow into concrete trenches

¹Southwest Fisheries Center Tiburon Laboratory, National Marine Fisheries Service, NOAA, Tiburon, CA 94920; present address: Auke Bay Fisheries Laboratory, National Marine Fisheries Service, NOAA, P.O. Box 155, Auke Bay, AK 99821.




Figure 1. — Schematic of Tiburon seawater system.




Figure 2. — Intake pumps, showing cleanouts and dual piping.



Figure 3. — Reservoir, aerator, brine tank, and pneumatic salt delivery line.



Figure 4.—Part of main wet laboratory. Plastic sheeting isolates tanks at rear for behavioral observations.

where it flows out a 6-inch cast iron line into the bay, or into 3-inch PVC recirculating pipes which deliver the water to the sump tank for recirculation (Fig. 5). Three-inch caps on Y fittings are located every few feet to allow entry to the recirculating lines.

All piping and valves in the system are schedule 80-PVC. Piping is in straight runs with threaded caps serving as cleanouts at each end. All seawater lines are dual to allow for maintenance, except for piping downstream from the UV sterilizer, where cleaning is minimal.

TEMPERATURE CONTROL

The seawater can be cooled to a set temperature ($\pm 2^\circ$ C). Although this temperature can be adjusted for the entire system, investigators must vary temperature in their tanks by individual cooling and heating devices.

A 750,000 BTU (British thermal unit) air-cooled compressor (Trane Model CGAA-6004-MA) cools ethylene glycol, which is circulated by one of two centrifugal pumps (Allis Chalmers Model F2L1, cast iron) through an all glass heat exchanger (Corning Model 600-GRB). An air-operated temperature probe in the seawater effluent



Figure 5.—Recirculating line in trench. Y fittings serve as entry points.

line from the heat exchanger controls the compressor through five steps of capacity. At the maximum step of capacity, 450 l/min can be cooled 10° C. The Corning heat exchanger is similar to that described by Lasker and Vlymen (1969).

SALINITY CONTROL

The salinity can be adjusted to any level above ambient ($\pm 0.5\text{‰}$). A 41,600-liter fiber glass brine tank is filled pneumatically with 31,745 kg of rock salt (Fig. 3). This salt is the unrefined product of a salt company located in the San Francisco Bay area. A typical analysis is given below:

Component	Percent
Calcium sulfate	0.09
Magnesium chloride	0.03
Magnesium sulfate	0.01
Moisture (H ₂ O)	3.45
Water Insolubles	0.01
Sodium chloride	96.41

Freshwater enters the tank, controlled by a float valve, and becomes saturated brine solution. This is metered into the seawater line by gravity flow through a stainless steel air-operated proportional valve, controlled by a conductivity meter pneumatic control system (Fig. 6). This system, patterned after Hettler et al. (1971), features conductivity meters with platinum probes mounted in the seawater line, a pneumatic controller, and chart recorder (Fig. 7). The temperature-compensated conductivity signal is converted to salinity, then recorded on a 7-day chart and fed into the pneumatic controller. The salinity is dialed on the controller, the instrument measures the difference between measured salinity and the set point, and sends a variable air pressure (4-15 pounds) to adjust the proportional valve. Variation in the salinity has been controlled to $\pm 0.5\text{‰}$. By adding another proportional valve connected to a freshwater source, one could lower the salinity if needed.

ALARM SYSTEM

An alarm has been incorporated into this system to avoid loss of fish and research time from component failures (Fig. 7). Specifically, the system alarms if: 1) water temperature exceeds high or low set points; 2) salinity exceeds high or low set points; 3) the water level is too high in sump tank; 4) the water level becomes too low in the reservoir; or 5) the electric power fails. An alarm signal activates a bell and timer. If the alarm persists past a preset time interval (30-300 s), a telephone dialer is activated, sending a signal to a local alarm company that has 24-h service (they have a list of people to call for corrective maintenance).

PERFORMANCE AND MAINTENANCE

The Tiburon Laboratory seawater system has been operating since April 1973. As of October 1974, after some modifications, the system is meeting the objectives of high water quality year-round.




Figure 6.—Schematic of salinity control system.

During September-December 1974, two filtered raw water samples and two recirculating water samples (entering the wet laboratories) were taken to determine water quality changes caused by recirculation with changing biomass. Samples were analyzed for nitrate, nitrite, and phosphate by a state-certified water quality laboratory. Temperature, pH, and salinity were measured at this facility (Table 1).

The results show no significant increase in nitrite, phosphate, and pH with recirculated water. Nitrate does accumulate at low levels but adjustment of the open flow will keep nutrients at acceptable levels. The December 9

condition indicates the seawater system water quality at near maximum biomass and flow rates.

At this biomass the oxygen content measured in the effluent of the tanks was 1-3 mg/l below saturation. The actual maximum biomass the system will maintain depends on the species held, temperature, and stress factors.

For the past year, although bay turbidities sometimes exceeded 100 Jackson Turbidity Units (JTU, Hach 2100 Turbidometer), values in the laboratories never exceeded 5 JTU. Because of this, cleaning of lines containing filtered water has been unnecessary. The absolute water clarity is important for the efficient functioning of the UV system (Herald et al. 1970).

The built-in UV output meter in our model (Refco Model EP-120, 316 stainless steel) indicates whether the unit needs cleaning or lamp replacement. To date, one cleaning has been needed (20% acetic acid soak for 1 h) and several lamps have needed replacement.

The temperature control has been less than satisfactory and will be modified. The high temperature variation of $\pm 2^\circ\text{C}$ is due to the cycling of the compressor through five stages of capacity to maintain the set temperature. A modification is planned whereby a two-way proportional valve controlled by an air-operated temperature controller will direct the ethylene glycol flow into the heat exchanger or a bypass line. This arrangement should allow for precise temperature control ($\pm 0.5^\circ\text{C}$). Plans also call for addition of heating capacity via a large-capacity water heater in the ethylene glycol line.

Salinity control has been precise ($\pm 0.5\text{‰}$) and has been maintained, although bay salinities varied from 12 to




Figure 7.—Control and alarm panel showing alarm indicators (top), conductivity meters (left), nullmatic controller (bottom center), and chart recorder (right).

Table 1. — Comparison of raw and recirculated water in the Tiburon seawater system.¹

Sample	Salinity (‰)	Temperature (°C)	pH	Total phosphate (mg/l)	Nitrate (mg/l)	Nitrite (mg/l)	Recirculating flow (l/min)	Open flow (l/min)	Biomass (kg)
Sept. 10									
Raw	26	18	7.8	0.80	0.44	0.02	300	60	20
Recirc.	26	16	7.8	0.62	0.70	0.02			
Oct. 29									
Raw	27	17	7.8	2.5	0.62	<0.01	240	55	30
Recirc.	26	17	7.8	2.8	0.61	<0.01			
Nov. 14									
Raw	23	13	7.8	0.5	0.31	<0.01	320	45	40
Recirc.	26	14	7.7	0.6	0.43	<0.01			
Dec. 9									
Raw	22	12	7.8	0.6	0.37	<0.01	400	200	340
Recirc.	24	11	7.8	0.6	0.52	0.02			

¹Mean values, $N = 2$. Total system volume = 65,000 liters.

30‰. Control can be maintained with up to 150 l/min open flow. With an average bay salinity of 18‰ and open circulation of 75 l/min, a level 10‰ higher can be maintained for \$270.00/week (salt cost \$30.00/ton, delivered). The cost of salt makes it mandatory to recirculate most of the flow and to monitor open circulation during periods of low bay salinities. It is recognized that the unnatural ionic balance of the salt used to increase salinity could cause problems with sensitive invertebrates or fish larvae, although no problems have been encountered to date.

No maintenance has been required except routine calibration. It is expected that the conductivity probes will need cleaning and replatinization every 2 yr, and the brine tank will need cleaning every 2-3 yr.

This seawater system has allowed us to hold and maintain a wide variety of organisms including adult and larval Pacific herring, adult and larval northern anchovy, juvenile striped bass, juvenile chinook salmon, and other bay fish and invertebrates. Organisms have been held for up to 8 mo and used for determining effects of petroleum


components on their physiology. Pacific herring and northern anchovy have been reared from eggs to juveniles. We have achieved our goal of year-round research with near-oceanic quality water.

ACKNOWLEDGMENTS

I thank Robert Brocksen, Reuben Lasker, and Jeanette Struhsaker for review of the manuscript.

LITERATURE CITED

- HERALD, E. S., R. P. DEMPSTER, and M. HUNT.
1970. Ultraviolet sterilization of aquarium water. *In* W. Hagen (editor), *Aquarium design criteria*. Special edition of *Drum and Croaker*, p. 57-71. U.S. Dep. Interior.
- HETTLER, W. F., R. W. LICHTENHELD, and H. R. GORDY.
1971. Open seawater system with controlled temperature and salinity. *Prog. Fish-Cult.* 33(1):3-11.
- LASKER, R., and L. L. VLYMEN.
1969. Experimental sea-water aquarium. *U.S. Fish Wildl. Serv., Circ.* 334, 14 p.



648. Weight loss of pond raised channel catfish (*Ictalurus punctatus*) during holding in processing plant vats. By Donald C. Greener and Robert J. Gill. December 1971. iv + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
649. Distribution of forage of skipjack tuna (*Katsuwonus pelamis*) in the eastern tropical Pacific. By Maurice Blackburn and Michael Myers. January 1972. iii + 16 p., 7 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
650. Effects of some antioxidants and EDTA in the development of rancidity in Spanish mackerel (*Scombrotoxa maculatus*) during frozen storage. By Robert N. Faraway. February 1972. iv + 12 p., 6 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
651. The effect of prenatal stress on drug metabolism in rainbow trout: the biochemistry and quality of skipjack tuna (*Katsuwonus pelamis*). April 1972. iii + 23 p., 3 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
652. The use of electricity in management of wild shrimp. Heavy production of Mexico shrimp trawl in Lake Michoacan. By Juan C. Ferris. March 1972. iv + 70 p., 11 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
653. An electric detector system for recognizing naturally oxygen-inflated fish genus *Brevoortia*. By R. O. Parker, Jr. February 1972. iv + 7 p., 1 fig., 1 appendix table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
654. Immobilization of energetic salmon and trout by decompression. By Doyle F. Sutherland. March 1972. iv + 7 p., 3 figs., 2 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
655. The calico scallop, *Argopecten gibber*. By Donald M. Aber and L. J. Costello. May 1972. iii + 19 p., 9 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
656. Making fish protein concentrates by enzymatic hydrolysis. A status report on research and some processes and products studied by NMF's. By Malcolm B. Hale. November 1972. iv + 32 p., 3 figs., 17 tables, 1 appendix table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
657. List of fishes of Alaska and adjacent waters with a guide to some of their literature. By Jay C. Quast and Elizabeth L. Heil. July 1972. iv + 47 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
658. The Southeast Fisheries Center biomerric code. Part 4. Fishes. By Harvey R. Bullis, Jr., Richard B. Roe, and Judith C. Gilbr. July 1972. vi + 95 p., 2 figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
659. A freshwater fish electro-motivator (FFEM) its characteristics and operation. By James E. Ellis and Charles C. Hoopes. November 1972. iii + 11 p., 9 figs.
660. A review of the literature on the development of skipjack tuna fisheries in the central and western Pacific Ocean. By Frank J. Hester and Tanno Otsu. January 1973. iii + 13 p., 1 fig. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
661. Seasonal distribution of tunas and billfishes in the Atlantic. By John P. Wise and Charles W. Davis. January 1973. iv + 24 p., 13 figs., 4 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
662. Fish larvae collected from the northeastern Pacific Ocean and Puget Sound during April and May 1967. By Kenneth D. Waldron. December 1972. iii + 16 p., 2 figs., 1 table, 4 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
663. Tagging and tag recovery experiments with Atlantic menhaden, *Brevoortia tyrannus*. By Richard L. Kroger and Robert L. Dryfoos. December 1972. iv + 11 p., 3 figs., 12 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
664. Larval fish survey of Humboldt Bay, California. By Maxwell B. Eldridge and Charles F. Bryan. December 1972. iii + 8 p., 8 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
665. Distribution of the common starbaster (*Homonotus noronhai*) in the lower Chesapeake Bay. August-August 1966 through December 1970. By James C. Quinn. August 1973. iv + 18 p., 18 figs., 11 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
666. An annotated bibliography of the cunner, *Tautoglabrus adspersus*. William T. Jones, M. Serchuk and David W. Frame. May 1973. iii + 43 p. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
667. A 14-day prediction for direct-readout meteorological satellites. By L. E. Anderson. August 1973. iv + 7 p., 2 figs., 1 table. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
668. Commercial fishery in the U.S. commercial fishery of western Lake Erie in 1969. By Thomas Van Meter. July 1973. iii + 11 p., 6 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
669. Coastal upwelling indices, west coast of North America, 1946-71. By Andrew Bakun. June 1973. iv + 103 p., 6 figs., 3 tables, 15 appendix figs. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
670. Seasonal occurrence of young Gull menhaden and other fishes in a northwestern Florida estuary. By Marian E. Tagatz and E. Peter H. Wilkins. August 1973. iii + 14 p., 1 fig., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
671. Abundance and distribution of inshore benthic fauna off southwestern Long Island, N.Y. By Frank W. Stemle, Jr. and Richard B. Stone. December 1973. iii + 50 p., 2 figs., 5 appendix tables.
672. Lake Erie bottom trawl explorations, 1962-66. By Edgar W. Bowman. January 1974. iv + 21 p., 9 figs., 1 table, 7 appendix tables.
673. Proceedings of the International Billfish Symposium, Kailua Kona, Hawaii, 9-12 August, 1972. Part 2. Review and Contributed Papers. Richard S. Shomura and Francis Williams (editors). July 1974. iv + 335 p., 38 papers. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
674. Price spreads and cost analyses for finfish and shellfish products at different marketing levels. By Erwin S. Penn. March 1974. vi + 74 p., 15 figs., 12 tables, 12 appendix figures, 11 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
675. Abundance of benthic macroinvertebrates in natural and altered estuarine areas. By Gill Gilmore and Lee Trent. April 1974. iii + 13 p., 11 figs., 3 tables, 2 appendix tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
676. Distribution, abundance, and growth of juvenile sockeye salmon, *Oncorhynchus nerka*, and associated species in the Naknek River system, 1961-64. By Robert J. Ellis. September 1974. v + 53 p., 27 figs., 26 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
677. Kinds and abundance of zooplankton collected by the USCG icebreaker *Glaciar* in the eastern Chukchi Sea, September-October 1970. By Bruce L. Wing. August 1974. iv + 18 p., 14 figs., 6 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
678. Pelagic amphipod crustaceans from the southeastern Bering Sea, June 1971. By Gerald A. Sanger. July 1974. iii + 8 p., 3 figs., 3 tables. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
679. Physiological response of the cunner, *Tautoglabrus adspersus*, to cadmium. October 1974. iv + 33 p., 6 papers, various authors. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 20402.
680. Heat exchange between ocean and atmosphere in the eastern North Pacific for 1961-71. By N. E. Clark, L. Eber, R. M. Laurs, J. A. Renner, and J. F. T. Saur. December 1974. iii + 108 p., 2 figs., 1 table, 5 plates.

UNITED STATES
DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL MARINE FISHERIES SERVICE
SCIENTIFIC PUBLICATIONS STAFF
ROOM 450
1107 N E 45TH ST
SEATTLE, WA 98105
OFFICIAL BUSINESS

POSTAGE AND FEES PAID
U S DEPARTMENT OF COMMERCE
COM-210



Marine Biological Laboratory S
Library - Periodicals
Woods Hole, Ma 02543



THIRD CLASS

